

AD-A152 821

CHEMICAL APPLICATIONS OF TOPOLOGY AND GROUP THEORY 17  
AN INFORMATION THEO. (U) GEORGIA UNIV ATHENS DEPT OF  
CHEMISTRY R B KING 27 MAR 85 TR-3 N00014-84-K-0365

1/1

UNCLASSIFIED

F/G 7/4

NL

END

FILMED

DTA



Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

2

## REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS  
BEFORE COMPLETING FORM

1. REPORT NUMBER Technical Report No. 3		2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) CHEMICAL APPLICATIONS OF TOPOLOGY AND GROUP THEORY. 17. AN INFORMATION THEORETICAL APPROACH TO POLYHEDRAL SYMMETRY		5. TYPE OF REPORT & PERIOD COVERED Technical Report	
7. AUTHOR(s) R. Bruce King		6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Georgia Department of Chemistry Athens, GA 30602		8. CONTRACT OR GRANT NUMBER(s) N00014-84-K-0365	
CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 051-861	
MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE March 27, 1985	
		13. NUMBER OF PAGES 17	
		15. SECURITY CLASS. (of this report)	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	

## DISTRIBUTION STATEMENT (of this Report)

This document has been approved for public release and sale; its distribution is unlimited.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		<b>DTIC</b> <b>ELECTE</b> <b>APR 25 1985</b>
18. SUPPLEMENTARY NOTES To be published in <u>Theoretica Chimica Acta</u>		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Polyhedra Symmetry Information Theory		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Information theoretic parameters are described which measure the asymmetry of polyhedra based on partitions of their vertices, faces, and edges into orbits under action of their symmetry point groups. Such asymmetry parameters are all zero only for the five regular polyhedra and are all unity for polyhedra having no symmetry at all, i.e., belonging to the $C_1$ symmetry point group. In all other cases such asymmetry parameters have values between zero and unity. Values for such asymmetry parameters are given for all topologically		

DTIC FILE COPY

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

85 04 01 172

AD-A152 821

## Block #20

distinct polyhedra having five, six, and seven vertices; all topologically distinct eight-vertex polyhedra having at least six symmetry elements; and selected polyhedra having from nine to twelve vertices. Effects of polyhedral distortions on these asymmetry parameters are examined for the tetrahedron, trigonal bipyramid, square pyramid, and octahedron. Such information theoretic asymmetry parameters can be used to order site partitions which are incomparable by the chirality algebra methods of Ruch and co-workers.

OFFICE OF NAVAL RESEARCH

Contract N00014-84-K-0365

Task No. NR 051-861

TECHNICAL REPORT NO. 3

Chemical Applications of Topology and Group Theory. 17. An Information

Theoretical Approach to Polyhedral Symmetry

by

R. Bruce King

Prepared for Publication

in

Theoretica Chimica Acta

University of Georgia  
Department of Chemistry  
Athens, Georgia

March 27, 1985

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Avail and/or	Special

A-1

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

This document has been approved for public release  
and sale; its distribution is unlimited.



# Abstract

C-sub1

Information theoretic parameters are described which measure the asymmetry of polyhedra based on partitions of their vertices, faces, and edges into orbits under action of their symmetry point groups. Such asymmetry parameters are all zero only for the five regular polyhedra and are all unity for polyhedra having no symmetry at all, i.e., belonging to the  $C_1$  symmetry point group. In all other cases such asymmetry parameters have values between zero and unity. Values for such asymmetry parameters are given for all topologically distinct polyhedra having five, six, and seven vertices; all topologically distinct eight-vertex polyhedra having at least six symmetry elements; and selected polyhedra having from nine to twelve vertices. Effects of polyhedral distortions on these asymmetry parameters are examined for the tetrahedron, trigonal bipyramid, square pyramid, and octahedron. Such information theoretic asymmetry parameters can be used to order site partitions which are incomparable by the chirality algebra methods of Ruch and co-workers.

## 1. Introduction

Symmetry is an important property of chemically significant polyhedra. In this connection a variety of descriptors can be used to define the symmetry of polyhedra. The most conventional polyhedral symmetry descriptor uses the symmetry point group [2]. Using this approach an increase in the symmetry of a polyhedron leads to an increase in the size of its point group. A related symmetry descriptor uses the cycle index polynomial for all of the symmetry operations of the polyhedron in question [3]. An increase in symmetry leads to more terms in the cycle index polynomial.

Such symmetry descriptors may be regarded as additive since an increase in symmetry leads to an increase in the size of the symmetry descriptor, i.e., the point group or the cycle index polynomial. Other alternative symmetry descriptors are subtractive. Chirality algebra [4,5,6] provides an example of a subtractive symmetry descriptor since an increase in the symmetry of the system decreases the number of chiral site partitions.

This paper discusses a new type of subtractive symmetry descriptor also based on site partitions but having information theory [7,8] rather than group representation theory [4,9] as its mathematical basis. This approach represents an extension of work of Bonchev, Kamenski, and Kamenska [8] on the information content of chemical structures. The approach in this paper defines information theoretical asymmetry parameters for the vertices, edges, and faces of a polyhedron such that these parameters are all zero for the five regular polyhedra [10] and all unity for polyhedra having no symmetry, i.e., polyhedra having

$C_1$  point group symmetry. These asymmetry parameters are functions solely of the site partitions of the vertices, the centers of the faces ("faces"), and the midpoints of the edges ("edges") of the polyhedron in question and in this sense have a similar genesis as the chirality functions [4,5] arising from chirality algebra. However, the fact that the asymmetry parameters are always fractions ranging from zero for systems in which all sites of a given type (i.e., vertices, faces, or edges) are equivalent (i.e., in the same orbit of the symmetry point group) to unity in systems having no symmetry (i.e., each site of a given type is its own orbit in the  $C_1$  point group) facilitates comparison of the symmetries of systems having radically different numbers of sites or symmetry point groups of different structures.

This paper defines such information theoretic asymmetry parameters for polyhedra. The values of these parameters are then examined for all polyhedra having seven or less vertices, all eight-vertex polyhedra having at least six symmetry elements, and selected polyhedra of chemical significance having nine through twelve vertices. Finally, this paper examines effects on such asymmetry parameters upon distortion of polyhedra of particular chemical importance: namely the tetrahedron, trigonal bipyramid, square pyramid, and octahedron.

## 2. Method

The polyhedron asymmetry parameters discussed in this paper are functions solely of the site partitions, where the sites are the vertices, the midpoints of the edges, or the midpoints of the faces. The site partitions are described by symbols of the type

$(a_1^{b_1} a_2^{b_2} \dots a_n^{b_n})$  where  $a_i$  and  $b_i$  are small positive integers and  $a_i \geq a_{i+1}$  ( $1 \leq i \leq n$ ). In this symbol for the site partition there are  $b_i$  sets of  $a_i$  identical sites. The  $a_i$  identical sites correspond to an orbit of the symmetry group. Thus, if all of the  $N$  sites of a given type (i.e., vertices, faces, or edges) are equivalent, the site partition is represented as  $(N^1)$ , abbreviated further as  $(N)$ . Conversely, if all of the  $N$  sites of a given type are different (i.e., if there is no symmetry whatsoever), the site partition is represented as  $(1^N)$ . For example, the site partitions of a trigonal bipyramid are  $(32)$  for the five vertices (i.e., three equatorial and two axial),  $(6)$  for the six (equivalent) faces, and  $(63)$  for the nine edges (i.e., six axial-equatorial and three equatorial-equatorial edges).

The information content of a site partition can be obtained from the following basic equation of Shannon<sup>7</sup>:

$$\bar{I} = - \sum_{i=1}^n p_i \lg p_i \quad (1)$$

In equation 1,  $n$  is the number of orbits,  $p_i$  is the probability of the site being in orbit  $i$ ,  $\lg$  is a logarithm to the base 2, and  $\bar{I}$  is the average information content per site. The probability  $p_i$  is obtained from the quotient  $N_i/N$  where  $N$  is the total number of sites and  $N_i$  is the number of sites in orbit  $i$ . For example, for the vertices of a trigonal bipyramid which correspond to a site partition  $(32)$ ,

$$\bar{I} = -(3/5)\lg(3/5) - (2/5)\lg(2/5) = 0.4422 + 0.5288 = 0.9710 \quad (2)$$

Note that if all of the sites are equivalent, there is only one orbit,

the probability of being in the orbit is 1 so that the average information content per site is zero, i.e.,  $\bar{I} = -\lg 1 = 0$ .

The maximum value of  $\bar{I}$  for a collection of  $N$  sites occurs when all sites are different, i.e., the system has no symmetry so that each site is its own orbit. For such a fully asymmetric system

$$\bar{I}^{\circ} = -\lg(1/N) \quad (3)$$

In equation 3  $\bar{I}^{\circ}$  represents the average information content per site for a fully asymmetric system. We can now define an asymmetry parameter  $A_s$  for  $N$  sites of type  $s$  (i.e., vertices, faces, or edges) by the quotient

$$A_s = \bar{I}/\bar{I}^{\circ} \quad (4)$$

where  $\bar{I}$  and  $\bar{I}^{\circ}$  are defined as in equations 1 and 3, respectively. For the vertices of a trigonal bipyramid with the site partition (32)

$$A_v(32) = \frac{-(3/5)\lg(3/5) - (2/5)\lg(2/5)}{-\lg(1/5)} = \frac{0.9710}{2.3221} = 0.4182 \quad (5)$$

Note that these asymmetry parameters depend only upon the site partitions. Furthermore, for  $N$  sites the asymmetry parameter for the fully symmetric site partition ( $N$ ) is 0, that for the fully asymmetric site partition ( $1^N$ ) is 1, and the asymmetry parameters for other site partitions fall between 0 and 1.

A further feature of the asymmetry parameter  $A_s$  defined in equation 4 is that for a given number of sites  $N$ ,  $A_s$  can only have a finite

number of discrete values, since there are only a relatively small number of ways for partitioning an integer  $N$  into a sum of smaller integers, i.e., 5, 7, 11, 14, and 22 such partitions for  $N = 4, 5, 6, 7$ , and 8, respectively. For this reason, only in a small number of exceptional cases other than the fully symmetric  $A(N) = 0$  and fully asymmetric  $A(1^N) = 1$  can asymmetry parameters be matched for partitions of different numbers of sites. These relatively rare matching of asymmetry parameters for small values of  $N$  include  $A(22) = A(422) = 0.5$  and  $A(21^2) = A(2^3 1^2) = 0.75$ .

A feature of the chirality algebra of Ruch and co-workers [4,5] is the recognition of some sets of different partitions of  $n$  sites as incomparable. This occurs when two different partitions of the same number of sites are ordered differently by different, equally valid, procedures. The simplest such pairs are the  $(3^2)$  and  $(41^2)$  partitions and the  $(2^3)$  and  $(31^3)$  partitions of six sites. The information theoretic asymmetry parameters for such incomparable site partitions may be distinct thereby providing a basis for ordering site partitions which are incomparable by the methods of Ruch and co-workers [4,5].

### 3. Results

The asymmetry parameters depend only on the site partitions and are given below for all possible partitions of four to eight sites:

A) Four Sites:  $A(4) = 0$ ;  $A(31) = 0.4057$ ;  $A(2^2) = 0.5$ ;  $A(21^2) = 0.75$ ;  $A(1^4) = 1$ .

B) Five Sites:  $A(5) = 0$ ;  $A(41) = 0.3109$ ;  $A(32) = 0.4182$ ;  $A(31^2) = 0.5905$ ;  $A(2^2 1) = 0.6555$ ;  $A(21^3) = 0.8278$ ;  $A(1^5) = 1$ .

C) Six Sites:  $A(6) = 0$ ;  $A(51) = 0.2515$ ;  $A(42) = 0.3552$ ;  $A(3^2) = 0.3868$ ;  
 $A(41^2) = 0.4842$ ;  $A(321) = 0.5645$ ;  $A(2^3) = 0.6132$ ;  $A(31^3) = 0.6935$ ;  
 $A(2^21^2) = 0.7421$ ;  $A(21^4) = 0.8711$ ;  $A(1^6) = 1$ .

D) Seven Sites:  $A(7) = 0$ ;  $A(61) = 0.2113$ ;  $A(52) = 0.3075$ ;  $A(43) = 0.3510$ ;  
 $A(51^2) = 0.4093$ ;  $A(421) = 0.4911$ ;  $A(3^21) = 0.5161$ ;  $A(32^2) = 0.5322$ ;  
 $A(41^3) = 0.5929$ ;  $A(321^2) = 0.6563$ ;  $A(31^4) = 0.7580$ ;  $A(2^21^3) = 0.7965$ ;  
 $A(21^5) = 0.8378$ ;  $A(1^7) = 1$ .

E) Eight Sites:  $A(8) = 0$ ;  $A(71) = 0.1812$ ;  $A(62) = 0.2704$ ;  $A(53) = 0.3182$ ;  
 $A(4^2) = 0.3333$ ;  $A(61^2) = 0.3537$ ;  $A(521) = 0.4329$ ;  $A(431) = 0.4686$ ;  
 $A(42^2) = 0.5$ ;  $A(51^3) = 0.5163$ ;  $A(3^22) = 0.5205$ ;  $A(421^2) = 0.5833$ ;  
 $A(3^21^2) = 0.6038$ ;  $A(32^21) = 0.6352$ ;  $A(41^4) = A(2^4) = 0.6667$ ;  
 $A(321^3) = 0.7186$ ;  $A(2^31^2) = 0.75$ ;  $A(31^5) = 0.8019$ ;  $A(2^21^4) = 0.8333$ ;  
 $A(21^6) = 0.9167$ ;  $A(1^8) = 1$ .

Table 1 lists the asymmetry parameters for all topologically distinct polyhedra having five, six and seven vertices. The properties of these polyhedra are taken from Federico's extensive tabulation of polyhedra having from four to eight faces [11] by conversion of the polyhedra to their duals [12,13]; the number of the dual of the polyhedron in question in Federico's table [11] is given to facilitate comparison. The polyhedra in Table 1 are ordered by increasing values of  $A_e$ , the edge asymmetry parameters, since among the three asymmetry parameters  $A_v$ ,  $A_f$ , and  $A_e$ , the parameter  $A_e$  has the maximum number of possible values because a given polyhedron has more edges than either vertices or faces by Euler's theorem, i.e.

$$v + f = e - 2$$

(6)

This, coupled with the intermediate dimensionality of edges (1) relative to vertices (0) and faces (2), suggests that  $A_e$  might be a better measure of polyhedral asymmetry than either  $A_v$  or  $A_f$ .

The asymmetry parameters of polyhedra having the common symmetry point groups fall into characteristic ranges. Thus the  $A_e$  values for polyhedra having the order 2 point groups  $C_s$  and  $C_2$  fall in the range 0.7 to 0.8 whereas those having the order 4 point group  $C_{2v}$  fall in the range 0.5 to 0.65. Furthermore, since the asymmetry parameters depend only on site partitions, all three asymmetry parameters will be identical for two or more polyhedra having identical site partitions for their vertices, faces, and edges. Such a set of polyhedra can be called isoentropic because of the relationship of information content to entropy [14]. Examples of isoentropic seven-vertex polyhedra include the seven seven-vertex polyhedra having no symmetry; a set of three seven-vertex polyhedra with  $A_v = 0.7964$ ,  $A_f = 0.7897$ , and  $A_e = 0.7749$ ; a set of three seven-vertex polyhedra with  $A_v = 0.7964$ ,  $A_f = 0.7500$ , and  $A_e = 0.7921$ ; and four pairs of isoentropic seven-vertex polyhedra having  $A_e$  values of 0.5578, 0.5943, 0.7506, and 0.7749 (Table 1). For a pair of dual [12,13] polyhedra P and P' (e.g., Federico dual numbers #35 and #38 in Table 1)  $A_e = A_e'$ ,  $A_v = A_f'$ , and  $A_f = A_v'$  in accord with the preservation of the symmetry of a polyhedron while constructing its dual.

According to Federico [11] the total number of combinatorically distinct eight-vertex polyhedra is 257, which is an intractable number for detailed study. However, if we exclude from consideration the large numbers of relatively uninteresting eight-vertex polyhedra having the relatively low symmetry point groups  $C_{2v}$ ,  $C_2$ ,  $C_s$ , and  $C_1$ , the

remaining number of eight-vertex polyhedra drops drastically to 14, a manageable number but still including the eight-vertex polyhedra of greatest chemical interest [15]. Table 2 lists the asymmetry parameters of some nine- to twelve-vertex polyhedra that have arisen in chemical contexts.

A given polyhedron has three asymmetry parameters  $A_v$ ,  $A_f$ , and  $A_e$  corresponding to the site partitions for the vertices, faces, and edges, respectively. All three of these parameters are zero only for the five regular polyhedra [10], namely the tetrahedron, octahedron (Table 1), cube (Table 2), icosahedron (Table 3), and regular (pentagonal) dodecahedron. Bipyramids, prisms, antiprisms, and the dual of the truncated tetrahedron (Table 2) have a single zero asymmetry parameter and the semiregular cuboctahedron [16] has zero values for  $A_v$  and  $A_e$  but not  $A_f$ .

Asymmetry parameters can also be used to follow the progress of distortion of relatively symmetrical polyhedra when symmetry elements are removed. Table 4 illustrates the effects of distortions on asymmetry parameters for four chemically significant polyhedra, namely the tetrahedron, trigonal bipyramid, square pyramid, and octahedron. Several different distortion pathways of the octahedron are examined in Table 4 depending on which symmetry elements (e.g., the  $C_3$  axis or a  $\sigma_h$  symmetry plane) are destroyed first in the distortion process. Note that as symmetry elements are removed in these distortion processes, the values of the asymmetry parameters increase in accord with expectations.

Acknowledgment: I am indebted to the U.S. Office of Naval Research for partial support of this work under Contract N00014-84-K-0365.

### References

1. For part 16 of this series see King, R.B.: Theoret. Chim. Acta (Berl.) 64, 453 (1984).
2. Cotton, F.A.: Chemical applications of group theory. New York: Wiley Interscience 1971.
3. King, R.B.: Inorg. Chem. 20, 363 (1981).
4. Ruch, E., Schönhofer, A.: Theoret. Chim. Acta (Berl.) 10, 91 (1968).
5. Ruch, E.: Accts. Chem. Res. 5, 49 (1983).
6. King, R.B.: Theoret. Chim. Acta (Berl.) 63, 103 (1983).
7. Shannon, C., Weaver, W.: Mathematical theory of communication. Urbana: University of Illinois 1949.
8. Bonchev, D., Kamenski, D., Kamenska, V.: Bull. Math. Biol. 38, 119 (1976).
9. Gorenstein, D.: Finite groups. New York: Harper and Row 1968.
10. Hilbert, D., Cohn-Vossen, S.: Geometry and the Imagination. New York: Chelsea 1952. pp. 89-93.
11. Federico, P.J.: Geom. Ded. 3, 469 (1975).
12. Grünbaum, B.: Convex polytopes. New York: Interscience, 1967.
13. King, R.B.: in Chemical applications of topology and graph theory, R.B. King, Ed. Amsterdam: Elsevier Scientific Publishing Company 1983.
14. Fast, J.D.: Entropy. Eindhoven, Holland: Philips 1968.
15. Kepert, D.L.: Prog. Inorg. Chem. 24, 179 (1978).
16. Lyusternik, L.A.: Convex Figures and Polyhedra. New York: Dover 1963. pp. 147-149.

TABLE 1

## ASYMMETRY PARAMETERS FOR TOPOLOGICALLY DISTINCT POLYHEDRA

HAVING FIVE, SIX, AND SEVEN VERTICES

Point Group	Partitions		Edges	Asymmetry Parameters			Federico Number of Dual	Comments
	Vertices	Faces		Vertices	Faces	Edges		
A) Five Vertex Polyhedra								
D <sub>3h</sub>	(32)	(6)	(63)	0.4182	0	0.2897	#2	trigonal bipyramid
C <sub>4v</sub>	(41)	(41)	(4 <sup>2</sup> )	0.3109	0.3109	0.3333	#3	square pyramid
B) Six Vertex Polyhedra								
O <sub>h</sub>	(6)	(8)	(12)	0	0	0	#7	octahedron
D <sub>3h</sub>	(6)	(32)	(63)	0	0.4182	0.2897	#10	trigonal prism
C <sub>5v</sub>	(51)	(51)	(5 <sup>2</sup> )	0.2515	0.2515	0.3010	#6	pentagonal pyramid
C <sub>2v</sub>	(2 <sup>3</sup> )	(421)	(4 <sup>2</sup> 21)	0.6132	0.4911	0.5270	#8	bicapped tetrahedron
C <sub>2v</sub>	(2 <sup>3</sup> )	(42 <sup>2</sup> )	(4 <sup>2</sup> 21 <sup>2</sup> )	0.6132	0.5000	0.5816	#4	
C <sub>2</sub>	(2 <sup>3</sup> )	(2 <sup>3</sup> )	(2 <sup>4</sup> 1 <sup>2</sup> )	0.6132	0.6132	0.7592	#9	
C <sub>s</sub>	(2 <sup>2</sup> 1 <sup>2</sup> )	(2 <sup>2</sup> 1 <sup>3</sup> )	(2 <sup>4</sup> 1 <sup>3</sup> )	0.7421	0.7964	0.7898	#5	
C) Seven Vertex Polyhedra								
D <sub>5h</sub>	(52)	(10)	(10,5)	0.3074	0	0.2350	#23	pentagonal bipyramid
C <sub>6v</sub>	(61)	(61)	(6 <sup>2</sup> )	0.2108	0.2108	0.2789	#19	hexagonal pyramid
C <sub>2v</sub>	(421)	(2 <sup>3</sup> )	(4 <sup>2</sup> 21)	0.4911	0.6132	0.5270	#44	

TABLE 1 (Continued)

Point Group	Partitions		Asymmetry Parameters			Federico Number of Dual	Comments	
	Vertices	Faces	Edges	Vertices	Faces			Edges
C <sub>3v</sub>	(3 <sup>2</sup> 1)	(3 <sup>2</sup> 1)	(3 <sup>4</sup> )	0.5161	0.5161	0.5578	#39, #40	Two isoentropic polyhedra
C <sub>2v</sub>	(421)	(42 <sup>2</sup> )	(4 <sup>2</sup> 2 <sup>2</sup> 1)	0.4911	0.5000	0.5843	#18	
C <sub>2v</sub>	(421)	(2 <sup>4</sup> )	(4 <sup>2</sup> 2 <sup>2</sup> 1)	0.4911	0.6667	0.5843	#36	
C <sub>2v</sub>	(421)	(42 <sup>2</sup> 1)	(4 <sup>2</sup> 2 <sup>3</sup> )	0.4911	0.5794	0.5872	#16	
C <sub>3v</sub>	(3 <sup>2</sup> 1)	(3 <sup>3</sup> 1)	(3 <sup>5</sup> )	0.5161	0.5706	0.5943	#12, #20	#20=capped octahedron
C <sub>2v</sub>	(2 <sup>3</sup> 1)	(42 <sup>2</sup> )	(4 <sup>2</sup> 2 <sup>3</sup> 1)	0.6947	0.4581	0.6246	#11	
C <sub>2</sub>	(2 <sup>3</sup> 1)	(2 <sup>3</sup> 1)	(2 <sup>6</sup> )	0.6947	0.6947	0.7211	#41	
C <sub>2</sub>	(2 <sup>3</sup> 1)	(2 <sup>4</sup> )	(2 <sup>6</sup> 1)	0.6947	0.6667	0.7506	#27, #37	
C <sub>8</sub>	(2 <sup>2</sup> 1 <sup>3</sup> )	(2 <sup>3</sup> 1 <sup>2</sup> )	(2 <sup>6</sup> 1)	0.7964	0.7500	0.7506	#31	
C <sub>2</sub>	(2 <sup>3</sup> 1)	(2 <sup>5</sup> )	(2 <sup>7</sup> 1)	0.6947	0.6990	0.7611	#13	
C <sub>8</sub>	(2 <sup>3</sup> 1)	(2 <sup>2</sup> 1 <sup>3</sup> )	(2 <sup>5</sup> 1 <sup>2</sup> )	0.6947	0.7964	0.7675	#38	#35 and #38 are dual
C <sub>8</sub>	(2 <sup>2</sup> 1 <sup>3</sup> )	(2 <sup>3</sup> 1)	(2 <sup>5</sup> 1 <sup>2</sup> )	0.7964	0.6947	0.7675	#35	#35 and #38 are dual
C <sub>2</sub> or C <sub>8</sub>	(2 <sup>2</sup> 1 <sup>3</sup> )	(2 <sup>4</sup> 1)	(2 <sup>6</sup> 1 <sup>2</sup> )	0.7964	0.7196	0.7749	#14, #24	Two isoentropic polyhedra
C <sub>8</sub>	(2 <sup>2</sup> 1 <sup>3</sup> )	(2 <sup>3</sup> 1 <sup>3</sup> )	(2 <sup>6</sup> 1 <sup>2</sup> )	0.7964	0.7897	0.7749	#21, #22, #28	Three isoentropic polyhedra
C <sub>8</sub>	(2 <sup>2</sup> 1 <sup>3</sup> )	(2 <sup>2</sup> 1 <sup>2</sup> )	(2 <sup>4</sup> 1 <sup>3</sup> )	0.7964	0.7421	0.7898	#43	
C <sub>8</sub>	(2 <sup>2</sup> 1 <sup>3</sup> )	(2 <sup>3</sup> 1 <sup>2</sup> )	(2 <sup>5</sup> 1 <sup>3</sup> )	0.7964	0.7500	0.7921	#17, #29, #30	Three isoentropic polyhedra
C <sub>1</sub>	(1 <sup>7</sup> )	(1 <sup>f</sup> )	(1 <sup>f</sup> +7-2) f=number of faces	1.0000	1.0000	1.0000	#15, #25, #26, #32, #33, #34, #42	Seven isoentropic polyhedra with no symmetry

f=number of faces

TABLE 2

ASYMMETRY PARAMETERS FOR TOPOLOGICALLY DISTINCT POLYHEDRA  
HAVING EIGHT VERTICES AND AT LEAST SIX SYMMETRY ELEMENTS

Point Group	Partitions		Edges	Asymmetry Parameters			Federico Number of Dual	Comments
	Vertices	Faces		Vertices	Faces	Edges		
$O_h$	(8)	(6)	(12)	0	0	0	#300	cube
$D_{6h}$	(62)	(12)	(12,6)	0.2704	0	0.2202	#54	hexagonal bipyramid
$T_d$	(4 <sup>2</sup> )	(12)	(12,6)	0.3333	0	0.2202	#49	dual of truncated tetrahedron
$D_{4d}$	(8)	(82)	(8 <sup>2</sup> )	0	0.2173	0.2500	#172	square antiprism
$C_{7v}$	(71)	(71)	(7 <sup>2</sup> )	0.1812	0.1812	0.2626	#247	heptagonal pyramid
$D_{2d}$	(4 <sup>2</sup> )	(4 <sup>2</sup> )	(842)	0.3333	0.3333	0.3621	#287, #288	two isoentropic polyhedra
$D_{2d}$	(4 <sup>2</sup> )	(82)	(84 <sup>2</sup> )	0.3333	0.2173	0.3750	#140	
$D_{3d}$	(62)	(6 <sup>2</sup> )	(6 <sup>3</sup> )	0.2704	0.2789	0.3801	#57	bicapped octahedron
$D_{3h}$	(62)	(63)	(6 <sup>2</sup> 3)	0.2704	0.2897	0.3895	#245	3,3-bicapped trigonal prism
$D_{2h}$	(4 <sup>2</sup> )	(4 <sup>2</sup> )	(82 <sup>3</sup> )	0.3333	0.3333	0.4372	#282	self-dual
$D_{2d}$	(4 <sup>2</sup> )	(84)	(84 <sup>2</sup> 2)	0.3333	0.2562	0.4404	#58	D <sub>2d</sub> -dodecahedron
$C_{3v}$	(3 <sup>2</sup> 12)	(63)	(63 <sup>3</sup> )	0.6038	0.2897	0.4419	#191	dual of triangular cupola
$C_{3v}$	(3 <sup>2</sup> 12)	(3 <sup>3</sup> )	(63 <sup>3</sup> )	0.6038	0.5000	0.4919	#194	

TABLE 3

## ASYMMETRY PARAMETERS FOR SELECTED POLYHEDRA HAVING

## NINE TO TWELVE VERTICES

Polyhedron	Point Group	Partitions			Asymmetry Parameters		
		Vertices	Faces	Edges	Vertices	Faces	Edges
Tricapped trigonal prism	D <sub>3h</sub>	(6,3)	(12,2)	(12,6,3)	0.2897	0.1554	0.3139
4-Capped square antiprism	C <sub>4v</sub>	(4 <sup>2</sup> 1)	(8,4,1)	(8,4 <sup>3</sup> )	0.4392	0.3348	0.4447
4,4-Bicapped square antiprism	D <sub>4d</sub>	(8,2)	(8 <sup>2</sup> )	(8 <sup>3</sup> )	0.2173	0.2500	0.3457
3,4,4,4,-Tetracapped trigonal prism	C <sub>3v</sub>	(3 <sup>3</sup> 1)	(12,3,1)	(12,3 <sup>4</sup> )	0.5706	0.2536	0.4362
B <sub>11</sub> H <sub>12</sub> - polyhedron	C <sub>2v</sub>	(4 <sup>2</sup> 3 <sup>1</sup> )	(4 <sup>3</sup> 2 <sup>3</sup> )	(4 <sup>5</sup> 2 <sup>3</sup> 1)	0.6321	0.6003	0.6417
Icosahedron	I <sub>h</sub>	(12)	(20)	(30)	0	0	0
Cuboctahedron	O <sub>h</sub>	(12)	(8,6)	(24)	0	0.2588	0

TABLE 4

## ASYMMETRY PARAMETERS OF DISTORTED POLYHEDRA

Point Group	Partitions		Asymmetry Parameters		Comments
	Vertices	Faces	Vertices	Faces	
A) Distorted Tetrahedra					
T <sub>d</sub>	(4)	(4)	(6)	0	regular tetrahedron
D <sub>2d</sub>	(4)	(4)	(42)	0	loss of C <sub>3</sub> axis
C <sub>3v</sub>	(31)	(31)	(32)	0.4057	trigonal pyramid
C <sub>2v</sub>	(22)	(22)	(412)	0.5000	loss of C <sub>3</sub> and S <sub>4</sub> axes
C <sub>s</sub>	(212)	(212)	(2212)	0.7500	reflection plane only
B) Distorted Trigonal Bipyramids					
D <sub>3h</sub>	(32)	(6)	(63)	0	maximum symmetry
C <sub>3v</sub>	(312)	(32)	(33)	0.5905	axial positions non-equivalent
C <sub>2v</sub>	(221)	(42)	(4221)	0.6554	loss of C <sub>3</sub> axis
C <sub>s</sub>	(213)	(2212)	(2313)	0.8277	reflection plane only
C) Distorted Square Pyramids					
C <sub>4v</sub>	(41)	(41)	(42)	0.3109	maximum symmetry: square base
C <sub>2v</sub>	(41)	(221)	(422)	0.3109	rectangular base
C <sub>2v</sub>	(221)	(41)	(422)	0.6554	rhombus base
C <sub>2</sub>	(221)	(221)	(24)	0.6554	parallelogram base
C <sub>s</sub>	(221)	(213)	(2312)	0.6554	trapezoid base

TABLE 4 (Continued)

Point Group	Partitions		Asymmetry Parameters			Comments	
	Vertices	Faces	Edges	Vertices	Faces		Edges
D) Distorted Octahedra							
$O_h$	(6)	(8)	(12)	0	0	0	regular octahedron
a) Three-fold Axis ( $C_3$ ) Retained							
$D_{3d}$	(6)	(62)	(6 <sup>2</sup> )	0	0.2704	0.2789	"trigonal antiprism"
$C_{3v}$	(3 <sup>2</sup> )	(3 <sup>2</sup> 1 <sup>2</sup> )	(63 <sup>2</sup> )	0.3868	0.6038	0.4184	no $\sigma_h$
b) Horizontal Symmetry Plane ( $\sigma_h$ ) Retained; no Three-fold Axis ( $C_3$ )							
$D_{4h}$	(42)	(8)	(84)	0.3552	0	0.2561	square cross-section
$D_{2h}$	(42)	(4 <sup>2</sup> )	(82 <sup>2</sup> )	0.3552	0.3333	0.3491	rectangle cross-section
$D_{2h}$	(2 <sup>3</sup> )	(8)	(4 <sup>3</sup> )	0.6132	0	0.4421	rhombus cross-section
$D_2$	(2 <sup>3</sup> )	(4 <sup>2</sup> )	(4 <sup>2</sup> 2 <sup>2</sup> )	0.6132	0.3333	0.5351	parallelogram cross-section
$C_{2v}$	(2 <sup>3</sup> )	(422)	(4 <sup>2</sup> 21 <sup>2</sup> )	0.6132	0.5000	0.5741	trapezoid cross-section
c) No Horizontal Symmetry Plane ( $\sigma_h$ ) or Three-fold Axis ( $C_3$ )							
$C_{4v}$	(41 <sup>2</sup> )	(4 <sup>2</sup> )	(4 <sup>3</sup> )	0.4842	0.3333	0.4421	square cross-section
$C_{2v}$	(41 <sup>2</sup> )	(2 <sup>4</sup> )	(4 <sup>2</sup> 2 <sup>2</sup> )	0.4842	0.6667	0.5351	rectangle cross-section
$C_{2v}$	(2 <sup>2</sup> 1 <sup>2</sup> )	(4 <sup>2</sup> )	(42 <sup>4</sup> )	0.7421	0.3333	0.6281	rhombus cross-section
$C_2$	(2 <sup>2</sup> 1 <sup>2</sup> )	(2 <sup>4</sup> )	(2 <sup>6</sup> )	0.7421	0.6667	0.7211	parallelogram cross-section
$C_s$	(2 <sup>2</sup> 1 <sup>2</sup> )	(2 <sup>2</sup> 1 <sup>4</sup> )	(2 <sup>5</sup> 1 <sup>2</sup> )	0.7421	0.8333	0.7676	trapezoid cross-section

**END**

**FILMED**

**5-85**

**DTIC**